

# A Diffeomorphic Positive Polynomial Map and its Implications in Chemistry

Gilles Gnacadja

Research and Development Information Systems  
Amgen

# Outline

- 1 Global Univalence
  - Background Material
  - A Diffeomorphic Positive Polynomial Map
- 2 Numerical Solution of the Polynomial Equation
  - A Particular Case
  - The General Case
- 3 Chemical Networks of Reversible Binding Reactions
  - Introduction to Complete Networks
  - Equilibrium State of Complete Networks
- 4 Closing Remarks

# Outline

- 1 Global Univalence
  - Background Material
  - A Diffeomorphic Positive Polynomial Map
- 2 Numerical Solution of the Polynomial Equation
  - A Particular Case
  - The General Case
- 3 Chemical Networks of Reversible Binding Reactions
  - Introduction to Complete Networks
  - Equilibrium State of Complete Networks
- 4 Closing Remarks

## Injectivity in Dimension One vs. in Dimension Two or Higher

## Injectivity in Dimension One

Let  $D$  be an interval in  $\mathbb{R}$  and  $f : D \rightarrow \mathbb{R}$  be a differentiable map.

$$f'(x_0) \neq 0 \quad \Rightarrow \quad \text{Local injectivity at } x_0 \quad (\text{Inv Fct Thm})$$

$$\forall x \in D, f'(x) \neq 0 \quad \Rightarrow \quad \text{Global injectivity on } D \quad (\text{Mean Val Thm})$$

## Injectivity in Dimension Two or Higher

Let  $D$  be an interval in  $\mathbb{R}^n$  and  $f : D \rightarrow \mathbb{R}^n$  be a differentiable map.

$$\det(f'(x_0)) \neq 0 \quad \Rightarrow \quad \text{Local injectivity at } x_0 \quad (\text{Inv Fct Thm})$$

$$\forall x \in D, \det(f'(x)) \neq 0 \quad \not\Rightarrow \quad \text{Global injectivity on } D$$

## Non-Injective Map in Dim Two with Positive Jacobian Determinant

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, x = (x_1, x_2) \mapsto f(x) = (e^{x_1} \cos x_2, e^{x_1} \sin x_2)$$

$$f'(x) = \begin{pmatrix} e^{x_1} \cos x_2 & -e^{x_1} \sin x_2 \\ e^{x_1} \sin x_2 & e^{x_1} \cos x_2 \end{pmatrix} \quad \det(f'(x)) = e^{2x_1} > 0$$

$f$  is constant on  $\{x_1\} \times (x_2 + 2\pi\mathbb{Z})$  for every  $(x_1, x_2) \in \mathbb{R}^2$ .

# P-matrix - Definition and Examples

## Definition

A real square matrix is a P-matrix (resp. a  $P_0$ -matrix) if all its principal minors are positive (resp. nonnegative).

## Examples

- A real positive-definite (resp. positive-semidefinite) matrix is a P-matrix (resp. a  $P_0$ -matrix).
- Let  $D$  and  $M$  be real  $n \times n$  matrices with  $D = \text{diag}(d_1, \dots, d_n)$ .
  - If  $d_1, \dots, d_n \geq 0$  and  $M$  is a  $P_0$ -matrix, then  $D + M$  is  $P_0$ -matrix.
  - If  $d_1, \dots, d_n \geq 0$  and  $M$  is a P-matrix, or if  $d_1, \dots, d_n > 0$  and  $M$  is a  $P_0$ -matrix, then  $D + M$  is P-matrix.

# P-matrix - Geometric Characterization

## Sign Reversal - Definition and Example

Let  $A$  be a real square matrix, let  $x$  be a real  $n$ -vector, and  $y = Ax$ . The matrix  $A$  reverses the sign of  $x$  if  $x_i y_i \leq 0$  for all  $i = 1, \dots, n$ .

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \quad u = \begin{pmatrix} -5 \\ 3 \end{pmatrix} \quad v = \begin{pmatrix} 2 \\ -1 \end{pmatrix} \quad w = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

$$Au = \begin{pmatrix} 1 \\ -3 \end{pmatrix} \quad Av = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \quad Aw = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Matrix  $A$  reverses the sign of vectors  $u$  and  $v$  but not of vector  $w$ .

## Theorem Geometric Characterization of a P-matrix

A real square matrix is a P-matrix if and only if it does not reverse the sign of any nonzero vector.

# P-Matrices and Global Univalence

## Global Univalence Theorem of Gale and Nikaidô

Let  $D$  be a rectangle in  $\mathbb{R}^n$  and let  $f : D \rightarrow \mathbb{R}^n$  be a differentiable function. If for every  $x \in D$ , the Jacobian matrix  $J(f, x)$  of  $f$  at  $x$  is a P-matrix, then  $f$  is injective.

GALE, D. and H. NIKAIÐÔ  
Math. Annalen 159, 81—93 (1965)

### **The Jacobian Matrix and Global Univalence of Mappings\***

By

DAVID GALE and HUKUKANE NIKAIÐÔ in Providence and Osaka

<http://dx.doi.org/10.1007/BF01360282>

(Theorem 4 and Remark 4.3)

Proof exploits features of the geometric characterization of P-matrices.

# Terminology: Univalence vs. Injectivity

- “Univalent” = “(Globally) Injective” = “One-to-One”.
- *Speculative thoughts:*

It appears that mathematical economists were originally more concerned with global injectivity than mainstream mathematicians. They had to address questions of uniqueness of equilibrium in various economic models. They made major contributions to the field and probably set the terminology.
- The condition “all principal minors are positive” that defines P-matrices is known in economics as the Hawkins-Simon condition after David Hawkins and Herbert A. Simon. Simon received the 1978 Nobel Prize in Economics (not for P-matrices).



# A Diffeomorphic Positive Polynomial Map

## Theorem

Let  $I$  be a finite subset of  $\mathbb{Z}_{\geq 0}^n$ , let  $(a_\alpha)_{\alpha \in I}$  be a family over  $\mathbb{R}_{\geq 0}$ , and let the function  $f = (f_1, \dots, f_n) : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}^n$  be given by

$$f_i(x) = x_i + \sum_{\alpha \in I} \alpha_i a_\alpha x^\alpha.$$

The function  $f$  is a  $C^\infty$ -diffeomorphism.

Notation:  $x^\alpha = (x_1, \dots, x_n)^{(\alpha_1, \dots, \alpha_n)} = x_1^{\alpha_1} \dots x_n^{\alpha_n}$ .

# Proof of Injectivity

- $f_i(x) = x_i + \sum_{\alpha \in I} \alpha_i a_\alpha x^\alpha$ .
- $J(f, x) = \text{Id}_n + M(x)$  with  $M(x) \cdot \text{diag}(x) = \sum_{\alpha \in I} a_\alpha x^\alpha \alpha^T \alpha$ .
- Matrix  $\alpha^T \alpha = (\alpha_i \alpha_j)_{1 \leq i, j \leq n}$  is positive-semidefinite.
- $M(x) \cdot \text{diag}(x)$  is positive-semidefinite.
- $M(x) \cdot \text{diag}(x)$  is a  $P_0$ -matrix.
- $M(x)$  is a  $P_0$ -matrix if  $x_1, \dots, x_n > 0$ .
- $M(x)$  is a  $P_0$ -matrix if  $x_1, \dots, x_n \geq 0$ .
- $J(f, x)$  is a P-matrix.
- Theorem of Gale and Nikaidô:  $f$  is injective.

# Proof of Local Diffeomorphism

- $f_i(\mathbf{x}) = x_i + \sum_{\alpha \in I} \alpha_i a_\alpha \mathbf{x}^\alpha$ .
- $f$  is defined and infinitely differentiable on  $\mathbb{R}^n$ .
- $\Omega := \{\mathbf{x} \in \mathbb{R}^n : \det(J(f, \mathbf{x})) > 0\}$ .
- $\Omega \supset \mathbb{R}_{\geq 0}^n$ .
- $\Omega$  is an open set.
- Inverse Function Theorem:  
 $f$  is a local  $C^\infty$ -diffeomorphism throughout  $\Omega$ .

# Proof of Surjectivity

- $f_i(x) = x_i + \sum_{\alpha \in I} \alpha_i a_\alpha x^\alpha$ .
- $F = (F_1, \dots, F_n) : \mathbb{R}_{\geq 0}^n \times \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}^n$   

$$F_i(b, x) = \frac{b_i}{1 + \sum_{\alpha \in I, \alpha_i \geq 1} \alpha_i a_\alpha x^{\alpha - e_{n,i}}}.$$
- $f^{-1}(b) = \text{FixPt}(F(b, -))$ .
- $F(b, \mathbb{R}_{\geq 0}^n) \subseteq [0_n, b]$ .
- Brouwer Fixed-Point Theorem:  $\text{FixPt}(F(b, -)) \neq \emptyset$ .

# Outline

- 1 Global Univalence
  - Background Material
  - A Diffeomorphic Positive Polynomial Map
- 2 Numerical Solution of the Polynomial Equation
  - A Particular Case
  - The General Case
- 3 Chemical Networks of Reversible Binding Reactions
  - Introduction to Complete Networks
  - Equilibrium State of Complete Networks
- 4 Closing Remarks

## Algorithm for a Particular Case

## Theorem

Let  $A = (a_{ij})_{1 \leq i, j \leq n}$  with  $a_{ij} \in \mathbb{R}_{\geq 0}$ , and  $b = (b_1, \dots, b_n) \in \mathbb{R}_{\geq 0}^n$ .  
The equation

$$x_i + \sum_{j=1}^n a_{ij} x_i x_j = b_i, \quad i = 1, \dots, n$$

has a unique solution  $x = (x_1, \dots, x_n) \in \mathbb{R}_{\geq 0}^n$ .

Let  $F = (F_1, \dots, F_n) : \mathbb{R}_{\geq 0}^n \times \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}^n$  with

$$F_i(b, x) = \frac{b_i}{1 + \sum_{j=1}^n a_{ij} x_j}.$$

Iterations of  $F(b, -)$  from  $b$  produce a sequence that encloses and linearly converges to the solution.

# Partial Proof with Diffeomorphic Polynomial Map

If the matrix  $A$  is symmetric, then the existence and uniqueness of the solution is a special case of the general diffeomorphism with

$$\begin{aligned} I &= \{ \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n : \|\alpha\|_{\ell^1} = \alpha_1 + \dots + \alpha_n = 2 \} \\ &= \{ 2e_{n,i} : 1 \leq i \leq n \} \cup \{ e_{n,ij} : 1 \leq i < j \leq n \} \end{aligned}$$

and

$$a_{2e_{n,i}} = a_{ij}/2$$

$$a_{e_{n,ij}} = a_{ij} .$$

# Complete Proof with Banach Contraction Theorem

## Theorem

With respect to the Thompson metric, the map  $F(b, -)$  is a contraction with Lipschitz constant  $\leq k = \frac{\|Ab\|_\infty}{1 + \|Ab\|_\infty}$ .

MATHEMATICAL METHODS IN THE APPLIED SCIENCES

*Math. Meth. Appl. Sci.* 2007; 30:201–211

Published online 23 October 2006 in Wiley InterScience

(www.interscience.wiley.com) DOI: 10.1002/mma.782

MOS subject classification: 47 H 10; 74 G 25; 74 G 30; 74 G 15; 92 C 40



Fixed points of order-reversing maps in  $\mathbb{R}_{>0}^n$   
and chemical equilibrium

Gilles Gnacadja\*†

*Systems Informatics Department, Amgen, Inc., One Amgen Center Drive, Thousand Oaks,  
California 91320-1799, U.S.A.*

Communicated by A. Kirsch

<http://dx.doi.org/10.1002/mma.782>  
(Theorem 5.4)



# Algorithms for General Case

**Wanted:** Algorithms for general case with *a priori* assurance of convergence and computational efficiency and accuracy.


**Project:** Consider interval methods.

# Outline

- 1 Global Univalence
  - Background Material
  - A Diffeomorphic Positive Polynomial Map
- 2 Numerical Solution of the Polynomial Equation
  - A Particular Case
  - The General Case
- 3 Chemical Networks of Reversible Binding Reactions
  - Introduction to Complete Networks
  - Equilibrium State of Complete Networks
- 4 Closing Remarks

## Two Networks of Actual Pharmacological Relevance



$$+ \quad +$$


$$+ \quad +$$


# The Problem

- Questions about equilibrium state:
  - Existence
  - Uniqueness, and with respect to what
  - Whether detailed-balanced
  - Asymptotic stability
  - Computation
  - Monotonicity with respect to particular species
  - etc.
- Observations:
  - Questions are addressed on a case-by-case basis.
  - Calculations are done with approximations that even some non-mathematicians bioscientists have questioned.
- Our effort: Devise a framework to encompass networks of interest and systematically address these questions.

# Complete Networks – The Idea

## Structure

- Reversible binding reactions:  
(*Two or more species*)  $\rightleftharpoons$  (*One species*).
- Notions of elementary species and composite species.  
Composite : Target/source of a binding/dissociation reaction.  
Elementary : Not composite.
- Notions of composition with respect to elementary species.
- Composition identifies species.
- Reactions preserve composition.

## Kinetics

- Law of Mass Action.
- Rate constants yield “consistent” cooperativity factors.

# Complete Networks – Realizing the Idea

Let  $I$  finite  $\subset \mathbb{Z}_{\geq 0}^n \setminus \{\|\cdot\|_{\ell^1} \leq 1\}$  and let  $(J_\alpha)_{\alpha \in I}$  be a family of finite, nonempty, pairwise disjoint subsets of  $\mathbb{Z}_{\geq 0}^{[1..n] \sqcup I} \setminus \{\|\cdot\|_{\ell^1} \leq 1\}$ .

## Complete Network of Reversible Binding Reactions

- **Elementary species** :  $X_1, \dots, X_n$
- **Composite species** :  $Y_\alpha, \alpha \in I$  ( $Y_\alpha = (X_1)_{\alpha_1} \cdots (X_n)_{\alpha_n}$ )
- **Reactions** :  $\sum_{i=1}^n \sigma_i X_i + \sum_{\beta \in I} \sigma_\beta Y_\beta \rightleftharpoons Y_\alpha, \alpha \in I, \sigma \in J_\alpha$   
 Kinetic : Law of Mass Action  
 $k_{\alpha,\sigma}^+, k_{\alpha,\sigma}^-$  : Association, dissociation rate constants  
 $a_{\alpha,\sigma} = k_{\alpha,\sigma}^+ / k_{\alpha,\sigma}^-$  : Equilibrium constants
- **Conservation of composition** :  
 $\alpha = (\sigma_1, \dots, \sigma_n) + \sum_{\beta \in I} \sigma_\beta \beta, \forall \alpha \in I, \forall \sigma \in J_\alpha$
- **Principle of detailed balance** : There exists a family  $(a_\alpha)_{\alpha \in I}$  over  $\mathbb{R}_{>0}$  such that  $a_\alpha = a_{\alpha,\sigma} \prod_{\beta \in I} (a_\beta)^{\sigma_\beta}, \forall \alpha \in I, \forall \sigma \in J_\alpha$ .



## Examples Revisited



$$n = 4$$

$$l: \begin{pmatrix} 1, 1, 0, 0 \\ 0, 1, 1, 0 \\ 0, 0, 1, 1 \\ 1, 0, 0, 1 \end{pmatrix}$$

Detailed-balance condition: Void.



$$n = 3$$

$$l: \begin{pmatrix} 1, 1, 0 \\ 1, 0, 1 \\ 1, 1, 1 \end{pmatrix}$$

Detailed-balance condition:

$$\begin{aligned} & a_{(R+A \rightleftharpoons RA)} a_{(RA+B \rightleftharpoons RAB)} \\ &= a_{(R+B \rightleftharpoons RB)} a_{(RB+A \rightleftharpoons RAB)} \cdot \end{aligned}$$

## Total Concentrations of Elementary Species

### Definition/Theorem

The total concentration of free and bound elementary species  $X_i$  is

$$[X_i] + \sum_{\alpha \in I} \alpha_i [Y_\alpha]$$

and is independent of time.

Notation:  $[\cdot]$  = concentration.



## Equil.: Existence - Uniqueness - Characterization - Stability

## Theorem

Let  $b = (b_1, \dots, b_n) \in \mathbb{R}_{\geq 0}^n$ .

- The complete network has a unique equilibrium state at which the total concentrations of elementary species  $X_1, \dots, X_n$  are  $b_1, \dots, b_n$ . Let  $x_i$  and  $y_\alpha$  denote the concentrations of  $X_i$  and  $Y_\alpha$  at that state.
- The vector  $x = (x_1, \dots, x_n)$  is the unique solution in  $\mathbb{R}_{\geq 0}^n$  of the equation  $x_i + \sum_{\alpha \in I} \alpha_i a_\alpha x^\alpha = b_i, i = 1, \dots, n$ .
- $y_\alpha = a_\alpha x^\alpha, \alpha \in I$ .
- The equilibrium state is detailed-balanced.
- If a concentrations trajectory originates at a state where the total concentrations of elementary species  $X_1, \dots, X_n$  are  $b_1, \dots, b_n$ , then it converges to this equilibrium state.

# Outline

- 1 Global Univalence
  - Background Material
  - A Diffeomorphic Positive Polynomial Map
- 2 Numerical Solution of the Polynomial Equation
  - A Particular Case
  - The General Case
- 3 Chemical Networks of Reversible Binding Reactions
  - Introduction to Complete Networks
  - Equilibrium State of Complete Networks
- 4 Closing Remarks

## Potential Future Work

- Enlarge the class of networks.
  - Species not necessarily determined by composition alone.  
Allow isomers.
  - Network not necessarily reversible.  
Allow weakly reversible, i.e. strongly connected.
  - Reactions not just binding and dissociation.  
Allow multiple species at both ends of reactions.
- Algorithms to solve the polynomial equation in all cases.
- Other uses for the diffeomorphic positive polynomial map.  
Applicable models in economics or other fields?